						—,			—,	—			—,	—,				
		/.	/ar	15	/is	/.	/ ,		/30	13		/5	/.	100	/20	/ à	/,	/x/
		ustralia P	ighanistan	andadest	ambodia	Chino/	DRE /	India	ndonesia	Mannat	Nepal	akista	PHG S	niippines	i lank	hailatt	Taiwar	Jiet Nam
	/ 4	77/ 4	%/ ¢		"/			/ `		*/	/		/ <	~	"/	`/	/`	·/
Infant Feeding Practices/Current Situation (outputs)	1	ĺ																
Low EBF rates in non-emergency times																		
High rates of bottle/formula feeding in non-emergency times																		
3. General perception that formula is as good as or better than breastmilk																		
MTCT of HIV an important cause of child mortality																		
Current IYCF/IFE Programme (inputs)																		
No national Code of Marketing or existing Code is not enforced																		
No government policy on IFE or policy is not implemented																		
7. Few government staffed trained on or knowledgeable about IFE																		
Few NGOs trained on or knowledgeable about IFE																		
Few lactation consultants during emergencies																		
10. Perception that women cannot breastfeeding during emergencies																		
11. Unsolicited donations of formula/milk powder received																		
12. Emergency food distributions do not contain suitable foods for CF																		
13. Emergency foods distributions often include formula or powdered milk																		
14. People in charge of emergency coordination do not prioritize IFE																		
NGOs operate independently of the national emergency response.																		
16. Rapid assessments seldom include information about IF																		
17. Poor understanding on indicators for IFE assessments																		
18. BF support services often not provided during emergencies																		
19. Emergency areas seldom have places suitable for women to BF																		
20. There are no systems to minimize risks of formula feeding																		